

Scientific Studies of the Effects of *Ionized Water*

Adequate fluid replacement helps maintain hydration and, promotes the health, safety, and optimal physical performance of individuals participating in regular physical activity.

Medical Science Sports Exercise. 1996 Jan;28(1):i-vii. American College of Sports Medicine position stand. Exercise and fluid replacement. Convertino VA, Armstrong LE, Coyle EF, Mack GW, Sawka MN, Senay LC Jr, Sherman WM.

It is the position of the American College of Sports Medicine that adequate fluid replacement helps maintain hydration and, therefore, promotes the health, safety, and optimal physical performance of individuals participating in regular physical activity. This position statement is based on a comprehensive review and interpretation of scientific literature concerning the influence of fluid replacement on exercise performance and the risk of thermal injury associated with dehydration and hyperthermia.

Based on available evidence, the American College of Sports Medicine makes the following general recommendations on the amount and composition of fluid that should be ingested in preparation for, during, and after exercise or athletic competition:

- 1) It is recommended that individuals consume a nutritionally balanced diet and drink adequate fluids during the 24-hr period before an event, especially during the period that includes the meal prior to exercise, to promote proper hydration before exercise or competition.
- 2) It is recommended that individuals drink about 500 ml (about 17 ounces) of fluid about 2 h before exercise to promote adequate hydration and allow time for excretion of excess ingested water.
- 3) During exercise, athletes should start drinking early and at regular intervals in an attempt to consume fluids at a rate sufficient to replace all the water lost through sweating (i.e., body weight loss), or consume the maximal amount that can be tolerated.
- 4) It is recommended that ingested fluids be cooler than ambient temperature [between 15 degrees and 22 degrees C (59 degrees and 72 degrees F)] and flavored to enhance palatability and promote fluid replacement. Fluids should be readily available and served in containers that allow adequate volumes to be ingested with ease and with minimal interruption of exercise.
- 5) Addition of proper amounts of carbohydrates and/or electrolytes to a fluid replacement solution is recommended for exercise events of duration greater than 1 h since it does not significantly impair water delivery to the body and may enhance performance. During exercise lasting less than 1 h, there is little evidence of physiological or physical performance differences between consuming a carbohydrate-electrolyte drink and plain water.
- 6) During intense exercise lasting longer than 1 h, it is recommended that carbohydrates be ingested at a rate of 30-60 g.h(-1) to maintain oxidation of carbohydrates and delay fatigue. This rate of carbohydrate intake can be achieved without compromising fluid delivery by drinking 600-1200 ml.h(-1) of solutions containing 4%-8% carbohydrates (g.100 ml(-1)). The carbohydrates can be sugars (glucose or sucrose) or starch (e.g., maltodextrin).
- 7) Inclusion of sodium (0.5-0.7 g.l(-1) of water) in the rehydration solution ingested during exercise lasting longer than 1 h is recommended since it may be advantageous in enhancing palatability, promoting fluid retention, and possibly preventing hyponatremia in certain individuals who drink excessive quantities of fluid. There is little physiological basis for the presence of sodium in oral rehydration solution for enhancing intestinal water absorption as long as sodium is sufficiently available from the previous meal.

Electrolyzed-reduced water scavenges active oxygen species and protects DNA from oxidative damage

Biochemical Biophysical Res Communication. 997 May 8;234(1):269-74. Shirahata S, Kabayama S, Nakano M, Miura T, Kusumoto K, Gotoh M, Hayashi H, Otsubo K, Morisawa S, Katakura Y. Institute of Cellular Regulation Technology, Graduate School of Genetic Resources Technology, Kyushu University, Fukuoka, Japan. sirahata@grt.kyushu-u.ac.jp

Active oxygen species or free radicals are considered to cause extensive oxidative damage to biological macromolecules, which brings about a variety of diseases as well as aging. The ideal scavenger for active oxygen should be 'active hydrogen'. 'Active hydrogen' can be produced in reduced water near the cathode during electrolysis of water. Reduced water exhibits high pH, low dissolved oxygen (DO), extremely high dissolved molecular hydrogen (DH), and extremely negative redox potential (RP) values. Strongly electrolyzed-reduced water, as well as ascorbic acid, (+)-catechin and tannic acid, completely scavenged O₂ produced by the hypoxanthine-xanthine oxidase (HX-XOD) system in sodium phosphate buffer (pH 7.0).

The superoxide dismutase (SOD)-like activity of reduced water is stable at 4 degrees C for over a month and was not lost even after neutralization, repeated freezing and melting, deflation with sonication, vigorous mixing, boiling, repeated filtration, or closed autoclaving, but was lost by opened autoclaving or by closed autoclaving in the presence of tungsten trioxide which efficiently adsorbs active atomic hydrogen. Water bubbled with hydrogen gas exhibited low DO, extremely high DH and extremely low RP values, as does reduced water, but it has no SOD-like activity. These results suggest that the SOD-like activity of reduced water is not due to the dissolved molecular hydrogen but due to the dissolved atomic hydrogen (active hydrogen). Although SOD accumulated H₂O₂ when added to the HX-XOD system, reduced water decreased the amount of H₂O₂ produced by XOD. Reduced water, as well as catalase and ascorbic acid, could directly scavenge H₂O₂. Reduced water suppresses single-strand breakage of DNA by active oxygen species produced by the Cu(II)-catalyzed oxidation of ascorbic acid in a dose-dependent manner, suggesting that reduced water can scavenge not only O₂·- and H₂O₂, but also 1O₂ and ·OH.

Comparison of electrolyzed oxidizing water with various antimicrobial interventions to reduce Salmonella species on poultry.

Poultry Science. 2002 Oct;81(10):1598-605. Fabrizio KA, Sharma RR, Demirci A, Cutter CN. Department of Food Science, The Pennsylvania State University, University Park 16802, USA.

Foodborne pathogens in cell suspensions or attached to surfaces can be reduced by electrolyzed oxidizing (EO) water; however, the use of EO water against pathogens associated with poultry has not been explored. In this study, acidic EO water [EO-A; pH 2.6, chlorine (CL) 20 to 50 ppm, and oxidation-reduction potential (ORP) of 1,150 mV], basic EO water (EO-B; pH 11.6, ORP of -795 mV), CL, ozonated water (OZ), acetic acid (AA), or trisodium phosphate (TSP) was applied to broiler carcasses inoculated with Salmonella Typhimurium (ST) and submerged (4 C, 45 min), spray-washed (85 psi, 25 C, 15 s), or subjected to multiple interventions (EO-B spray, immersed in EO-A; AA or TSP spray, immersed in CL). Remaining bacterial populations were determined and compared at Day 0 and 7 of aerobic, refrigerated storage. At Day 0, submersion in TSP and AA reduced ST 1.41 log₁₀, whereas EO-A water reduced ST approximately 0.86 log₁₀. After 7 d of storage, EO-A water, OZ, TSP, and AA reduced ST, with detection only after selective enrichment.

Spray-washing treatments with any of the compounds did not reduce ST at Day 0. After 7 d of storage, TSP, AA, and EO-A water reduced ST 2.17, 2.31, and 1.06 log₁₀, respectively. ST was reduced 2.11 log₁₀ immediately following the multiple interventions, 3.81 log₁₀ after 7 d of storage. Although effective against ST, TSP and AA are costly and adversely affect the environment. This study demonstrates that EO water can reduce ST on poultry surfaces following extended refrigerated storage.

Inactivation of Escherichia coli (O157:H7) and Listeria monocytogenes on plastic kitchen cutting boards by electrolyzed oxidizing water

Venkitanarayanan KS, Ezeike GO, Hung YC, Doyle MP. Department of Animal Science, University of Connecticut, Storrs 06269, USA.

One milliliter of culture containing a five-strain mixture of Escherichia coli O157:H7 (approximately 10¹⁰ CFU) was inoculated on a 100-cm² area marked on unscarred cutting boards. Following inoculation, the boards were air-dried under a laminar flow hood for 1 h, immersed in 2 liters of electrolyzed oxidizing water or sterile de-ionized water at 23 degrees C or 35 degrees C for 10 or 20 min; 45 degrees C for 5 or 10 min; or 55 degrees C for 5 min. After each temperature-time combination, the surviving population of the pathogen on cutting boards and in soaking water was determined.

Soaking of inoculated cutting boards in electrolyzed oxidizing water reduced E. coli O157:H7 populations by > or = 5.0 log CFU/100 cm² on cutting boards. However, immersion of cutting boards in de-ionized water decreased the pathogen count only by 1.0 to 1.5 log CFU/100 cm². Treatment of cutting boards inoculated with Listeria monocytogenes in electrolyzed oxidizing water at selected temperature-time combinations (23 degrees C for 20 min, 35 degrees C for 10 min, and 45 degrees C for 10 min) substantially reduced the populations of L. monocytogenes in comparison to the counts recovered from the boards immersed in de-ionized water. E. coli O157:H7 and L. monocytogenes were not detected in electrolyzed oxidizing water after soaking treatment, whereas the pathogens survived in the de-ionized water used for soaking the cutting boards. This study revealed that immersion of kitchen cutting boards in electrolyzed oxidizing water could be used as an effective method for inactivating food-borne pathogens on smooth, plastic cutting boards.

PMID: 10456736 [Pub Med - indexed for MEDLINE]

The bactericidal effects of electrolyzed oxidizing water on bacterial strains involved in hospital infections

Vorobjeva NV , Vorobjeva LI, Khodjaev EY. *Artificial Organs* 2004 Jun;28(6):590-2. Department of Physiology of Microorganisms, Biology Faculty, Moscow State University, Lenin Hills 1/12, Moscow 119992, Russia. nvvorobjeva@mail.ru

The study is designed to investigate bactericidal actions of electrolyzed oxidizing water on hospital infections. Ten of the most common opportunistic pathogens are used for this study. Cultures are inoculated in 4.5 mL of electrolyzed oxidizing (EO) water or 4.5 mL of sterile de-ionized water (control), and incubated for 0, 0.5, and 5 min at room temperature. At the exposure time of 30 s the EO water completely inactivates all of the bacterial strains, with the exception of vegetative cells and spores of bacilli which need 5 min to be killed. The results indicate that electrolyzed oxidizing water may be a useful disinfectant for hospital infections, but its clinical application has still to be evaluated. PMID: 15153153 [Pub Med - in process]

Effect of electrolyzed oxidizing water and hydrocolloid occlusive dressings on excised burn-wounds in rats

Chin J Traumatol. 2003 Aug 1;6(4):234-7. Xin H, Zheng YJ, Hajime N, Han ZG. Department of Thoracic Surgery , China - Japan Union Hospital , Jilin University, Jilin 130031, China . xinhua7254@yahoo.com.cn

OBJECTIVE: To study the efficacy of electrolyzed oxidizing water (EOW) and hydrocolloid occlusive dressings in the acceleration of epithelialization in excised burn-wounds in rats.

METHODS: Each of the anesthetized Sprague-Dawley rats (n=28) was subjected to a third-degree burn that covered approximately 10% of the total body surface area. Rats were assigned into four groups: Group I (no irrigation), Group II (irrigation with physiologic saline), Group III (irrigation with EOW) and Group IV (hydrocolloid occlusive dressing after EOW irrigation). Wounds were observed macroscopically until complete epithelialization was present, then the epithelialized wounds were examined microscopically.

RESULTS: Healing of the burn wounds was the fastest in Group IV treated with hydrocolloid occlusive dressing together with EOW. Although extensive regenerative epidermis was seen in each Group, the proliferations of lymphocytes and macrophages associated with dense collagen deposition were more extensive in Group II, III and IV than in Group I. These findings were particularly evident in Group III and IV.

CONCLUSIONS: Wound Healing may be accelerated by applying a hydrocolloid occlusive dressing on burn surfaces after they are cleaned with EOW.

PMID: 12857518 [Pub Med - indexed for MEDLINE]

Use of Ionized Water in hypochlorhydria or achlorhydria

Prof. Kuninaka Hironage, Head of Kuninaka Hospital

"Too many fats in the diets, which lead to the deposition of cholesterol on the blood vessels, which in turn constrict the blood flow, cause most illnesses such as high blood pressure. In accordance with the theory of Professor Gato of Kyushu University on Vitamin K (because vitamin K enables the blood calcium to increase) , or the consumption of more antioxidant water, the effectiveness of the increase in the calcium in high blood pressure is most significant. The consumption of alkaline antioxidant water for a period of 2 to 3 months, I have observed the blood pressure slowly drop, due to the water's solvent ability, which dissolves the cholesterol in the blood vessels ."

Effect of electrolyzed water on wound healing

Artif Organs. 2000 Dec;24(12):984-7. Yahagi N, Kono M, Kitahara M, Ohmura A, Sumita O, Hashimoto T, Hori K, Ning-Juan C, Woodson P, Kubota S, Murakami A, Takamoto S. Department of Anesthesiology, Teikyo University Mizonokuchi Hospital , Tokyo , Japan . naokiyah@aol.com

Electrolyzed water accelerated the healing of full-thickness cutaneous wounds in rats, but only anode chamber water (acid pH or neutralized) was effective. Hypochlorous acid (HOCl), also produced by electrolysis, was ineffective, suggesting that these types of electrolyzed water enhance wound healing by a mechanism unrelated to the well-known antibacterial action of HOCl. One possibility is that reactive oxygen species, shown to be electron spin resonance spectra present in anode chamber water, might trigger early wound healing through fibroblast migration and proliferation.

PMID: 11121980 [Pub Med - indexed for MEDLINE]

Physiological effects of alkaline Ionized Water: Effects on metabolites produced by intestinal fermentation by Takashi Hayakawa, Chicko Tushiya, Hisanori Onoda, Hisayo Ohkouchi, Harul--to Tsuge (Gifu University, Faculty of Engineering, Dept. of Food Science)

We have found that long-term ingestion of alkaline ionized water (AIW) reduces cecal¹ fermentation in rats that were given highly fermentable commercial diet (MF: Oriental Yeast Co., Ltd.). In this experiment, rats were fed MF and test water (tap water, AIW with pH at 9 and 10) for about 3 months. Feces were collected on the 57th day, and the rats were dissected on the 88th day. The amount of ammonium in fresh feces and cecal contents as well as fecal free-glucose tended to drop down for the AIW group. In most cases, the amount of free-amino acids in cecal contents did not differ significantly except for cysteine (decreased in AIW with pH at 10) and isoleucine (increased in AIW with pH at 10).

Purpose of tests

Alkaline ionized water electrolyzers have been approved for manufacturing in 1965 by the Ministry of Health and Welfare as medical equipment to produce medical substances. Alkaline ionized water (AIW) produced by this equipment is known to be effective against gastrointestinal fermentation, chronic diarrhea, indigestion and hyperchylia as well as for controlling gastric acid.*1 This is mainly based on efficacy of the official calcium hydroxide. *2 By giving AIW to rats for a comparatively long time under the condition of extremely high level of intestinal fermentation, we have demonstrated that AIW intake is effective for inhibition of intestinal fermentation when its level is high based on some test results where AIW worked against cecal hypertrophy and for reduction in the amount of short-chain fatty acid that is the main product of fermentation.*3 We have reported that this is caused by the synergy between calcium level generally contained in AIW (about 50ppm) and the value of pH, and that frequency of detecting some anaerobic bacteria tends to be higher in alkaline Ionized Water groups than the other, although the bacteria count in the intestine does not have significant difference.

Based on these results, we made a judgment that effect of taking AIW supports part of inhibition mechanism against abnormal intestinal fermentation, which is one of the claims of efficacy that have been attributed to alkaline ionized water electrolyzers. *4 On the other hand, under the dietary condition of low intestinal fermentation, AIW uptake does not seem to inhibit fermentation that leads us to believe that effect of AIW uptake is characteristic of hyper-fermentation state. Metabolites produced by intestinal fermentation include indole and skatole in addition to organic acids such as short-chain fatty acid and lactic acid as well as toxic metabolites such as ammonium, phenol and p-cresol. We do not know how AIW uptake would affect the production of these materials. In this experiment, we have tested on ammonium production as explained in the following sections.

Testing methods

Four-week-old male Wistar/ST Clean rats were purchased from Japan SLC Co., Ltd. and were divided into 3 groups of 8 each after preliminary breeding. AIW of pH 9 and 10 was produced by an electrolyzer Mineone ROYAL NDX3 1 OH by Omco Co., Ltd. This model produces AIW by electrolyzing water with calcium lactate added. On the last day of testing, the rats were dissected under Nembutal anesthesia to take blood from the heart by a heparin-treated syringe. As to their organs, the small intestines, cecum and colon plus rectum were taken out from each of them. The cecum was weighed and cleaned with physiological saline after its contents were removed, and the tissue weight was measured after wiping out moisture. Part of cecal contents was measured its pH, and the rest was used to assay ammonium concentration. The amount of ammonium contained in fresh feces and cecal contents was measured by the Nessler method after collecting it in the extracted samples using Conway's micro-diffusion container. Fecal free-glucose was assayed by the oxygen method after extraction by hot water. Analysis of free amino acids contained in cecal contents was conducted by the Waters PicoTag amino acid analysis system.

Test results and analyses

No difference was found in the rats' weight gain, water and feed intake and feeding efficiency, nor was any particular distinction in appearance identified. The length of the small intestines and colon plus rectum tended to decline in AIW groups. PH value of cecal contents was higher and the amount of fecal free-glucose tended to be lower in AIW groups than the control group. Since there was no difference in fecal discharge itself, the amount of free-glucose discharged per day was at a low level. The amount of discharged free-glucose in feces is greater when intestinal fermentation is more intensive, which indicates that intestinal fermentation is more inhibited in AIW groups than the control group. Ammonium concentration in cecal contents tends to drop down in AIW groups (Fig. 1). This trend was most distinctive in case of fresh feces of one of AIW groups with pH 10 (Fig.2) AIW uptake was found to be inhibitory against ammonium production. In order to study dynamics of amino acids in large intestines, we examined free amino acids in the cecal contents to find out that cysteine level is low in AIW groups whereas isoleucine level is high in one of AIW groups with pH 10, although no significant difference was identified for other amino acids.

Bibliography

"Verification of Alkaline Ionized Water" by Life Water Institute, Metamor Publishing Co., 1994, p.46

"Official Pharmaceutical Guidelines of Japan , Vol. IT" by Japan Public Documents Association, Hirokawa Publishing Co. , 1996

"Science and Technology of Functional Water" (part) by Takashi Hayakawa, Haruhito Tsuge, edited by Water Science Institute, 1999, pp.109-116

"Basics and Effective Use of Alkaline Ionized Water" by Takashi Hayakawa, Haruhito Tsuge, edited by Tetsuji Hekudou, 25th General Assembly of Japan Medical Congress Functional Water in Medical Treatment", Administration Offices, 1999, pp. 10- 11

Clinical Improvements Obtained From The Intake Of Reduced Water

Extracts from "Presentation At The Eight Annual International Symposium On man And His Environment in Health And Disease" on February 24th 1990, at The Grand Kempinski Hotel, Dallas, Texas, USA by Dr. H. Hayashi, M.D. and Dr. M Kawamura, M.D.,

Since the introduction of alkaline ionic water in our clinic in 1985, we have had the following interesting clinical experiences in the use of this type of water. By the use of alkaline ionic water for drinking and the preparation of meals for our in-patients, we have noticed :

- Declines in blood sugar levels in diabetic patients.
- Improvements in peripheral circulation in diabetic gangrene.
- Declines in uric acid levels in patients with gout.
- Improvements in liver function exams in hepatic disorders.
- Improvements in gastroduodenal ulcer and prevention of their recurrences.
- Improvements in hypertension and hypotension.
- Improvements in allergic disorders such as asthma, urticaria, rhinites and atopic dermatitis.
- Improvements in persistent diarrhea which occurred after gastrectomy.
- Quicker improvements in post operative bowel paralysis.
- Improvements in serum bilirubin levels in new born babies.

Being confirming clinical improvements, we have always observed changes of stools of the patients, with the color of their feces changing from black-brown color to a brighter yellow-brown one, and the odor of their feces becoming almost negligible.

The number of patients complaining of constipation also decreased markedly. The change of stool findings strongly suggests that alkaline ionic water intake can decrease the production of putrefied or pathogenic metabolites. Devices to produce reduced water were introduced into our clinic in May 1985. Based on the clinical experiences obtained in the past 15 years, it can be said that introduction of electrolyzed-reduced water for drinking and cooking purpose for in-patients should be the very prerequisite in our daily medical practices. Any dietary recipe cannot be a scientific one if property of water is not taken by the patients is not taken into consideration.

The Ministry of Health and Welfare in Japan announced in 1965 that the intake of reduced water is effective for restoration of intestinal flora metabolism.

Decomposition of ethylene, a flower-senescence hormone, with electrolyzed anode water. Biosci Biotechnology Biochemistry 2003 Apr;67(4):790-6. Harada K, Yasui K. Department of Research and Development, Hokkaido Electric Power Co., Inc., 2-1 Tsuishikari, Ebetsu, Hokkaido 067-0033, Japan. kharada@h1.hotcn.ne.jp

Electrolyzed anode water (EAW) markedly extended the vase life of cut carnation flowers. Therefore, a flower-senescence hormone involving ethylene decomposition by EAW with potassium chloride as an electrolyte was investigated. Ethylene was added externally to EAW, and the reaction between ethylene and the available chlorine in EAW was examined. EAW had a low pH value (2.5), a high concentration of dissolved oxygen, and extremely high redox potential (19.2 mg/l and 1323 mV, respectively) when available chlorine was at a concentration of about 620 microns.

The addition of ethylene to EAW led to ethylene decomposition, and an equimolar amount of ethylene chlorohydrine with available chlorine was produced. The ethylene chlorohydrine production was greatly affected by the pH value (pH 2.5, 5.0 and 10.0 were tested), and was faster in an acidic solution. Ethylene chlorohydrine was not produced after ethylene had been added to EAW at pH 2.6 when available chlorine was absent, but was produced after potassium hypochlorite had been added to such EAW. The effect of the pH value of EAW on the vase life of cut carnations was compatible with the decomposition rate of ethylene in EAW of the same pH value. These results suggest that the effect of EAW on the vase life of cut carnations was due to the decomposition of ethylene to ethylene chlorohydrine by chlorine from chlorine compounds.

PMID: 12784619 [Pub Med - indexed for MEDLINE]

Use of Ionized Water in treating Acidosis

Prof. Hatori Tasutaroo, Head of Akajiuji Blood Centre, Yokohama Hospital , Faitama District

" Due to a higher standard of living, our eating habits have changed. We consume too much proteins, fats and sugar. The excess fats and carbohydrates are in the body as fats. In the present lifestyles, Americans are more extravagant on food compared to the Japanese. Due to this excessive intake obesity is a significant problem. Normally, one out of five males and one out of four females is obese.

The degree of "burn-out" in food intake largely depends on the amount on intake of vitamins and minerals. When excessive intake of proteins, carbohydrates and fats occurs, the requirement for vitamins and minerals increases.

However, there is not much research carried out pertaining to the importance of vitamins and minerals.

Nowadays, many people suffer from acidification that leads to diabetes, heart diseases, cancer, liver and kidney diseases. If our food intake can be completely burned off, then there is no deposition of fats. Obviously, there will be no acidification problem and hence there should not be any sign of obesity.

The antioxidant water contains an abundance of ionic calcium. This ionic calcium helps in the "burn-off" process. By drinking antioxidant water, it provides sufficient minerals for our body. As a result, we do not need to watch our diet to stay slim.

Hence, antioxidant water is a savior for those suffering from obesity and many adult diseases, providing good assistance in enhancing good health."

Allergies and Ionized Water

Prof. Kuninaka Hironaga, Head of Kuninaka Hospital

"Mr. Yamada, the head of Police Research Institute, suffered from severe allergy. He was treated repeatedly by skin specialist, but with no success. Then he started consuming antioxidant water. The allergy responded very well and was soon completely cured. No relapse had occurred, although he had taken all kinds of food. He was most grateful and excited about this treatment.

As for myself, I had also suffered severe allergy. Ever since I began to consume antioxidant water, the allergy has recovered. Since then, I started a research on the effectiveness of antioxidant water.

I discovered that most allergies are due to acidification of body condition and is also related to consuming too much meat and sugar. In every allergy case, the patient's antioxidant minerals are excessively low which in turn lower the body resistance significantly. The body becomes overly sensitive and develops allergy easily. To stabilize the sensitivity, calcium solution is injected into the vein. Therefore, it is clear that the antioxidant water has ionic calcium, which can help alleviate allergy.

The ionic calcium not only enhances the heart, urination, and neutralization of toxins but controls acidity. It also enhances the digestive system and liver function. This will promote natural healing power and hence increase its resistance to allergy. In some special cases of illness, which do not respond to drugs, it is found, it is found to respond well to antioxidant water."

Use of Ionized Water for gynecological conditions and treatments

Prof. Watanabe Ifao, Watanabe Hospital

Ionized alkaline antioxidant water improves body constituents and ensures effective healing to many illnesses. The uses of antioxidant water in gynecological patients have proved to be very effective. The main reason for its effectiveness is that this water can neutralize toxins.

When given antioxidant water to pre-eclamptic toxemia cases, the results are most significant. During my long years of servicing the pre-eclamptic toxemia cases, I found that the women with pre-eclamptic toxemia who consumed antioxidant water tend to deliver healthier babies with stronger muscles . A survey report carried out on babies in this group showed intelligence above average.

The mechanism of the enhanced antioxidant effects against superoxide anion radicals of reduced water produced by electrolysis

Biophysical Chemistry. 2004 Jan 1;107(1):71-82. Hanaoka K, Sun D, Lawrence R, Kamitani Y, Fernandes G. Bio-REDOX Laboratory Inc. 1187-4, Oaza-Ueda, Ueda-shi, Nagano-ken 386-0001, Japan . hanak@rapid.ocn.ne.jp

We reported that reduced water produced by electrolysis enhanced the antioxidant effects of proton donors such as ascorbic acid (AsA) in a previous paper. We also demonstrated that reduced water produced by electrolysis of 2 mM NaCl solutions did not show antioxidant effects by itself. We reasoned that the enhancement of antioxidant effects may be due to the increase of the ionic product of water as solvent. The ionic product of water (pK_w) was estimated by

measurements of pH and by a neutralization titration method. As an indicator of oxidative damage, Reactive Oxygen Species- (ROS) mediated DNA strand breaks were measured by the conversion of supercoiled phiX-174 RF I double-strand DNA to open and linear forms. Reduced water had a tendency to suppress single-strand breakage of DNA induced by reactive oxygen species produced by H₂O₂/Cu (II) and HQ/Cu (II) systems. The enhancement of superoxide anion radical dismutation activity can be explained by changes in the ionic product of water in the reduced water.

Toxin Neutralization with the Use of Electrolysed Water

Prof. Kuwata Keijiroo, Doctor of Medicine

"In my opinion, the wonder of antioxidant water is the ability neutralizes toxins, but it is not a medicine. The difference is that the medicine can only apply to each and individual case, whereas the antioxidant water can be consumed generally and its neutralizing power is something which is very much unexpected. Now, in brief, let me introduce to you a heart disease case and how it was cured.

The patient was a 35 years old male suffering from vascular heart disease. For 5 years, his sickness deteriorated. He was in the Setagays Government Hospital for treatment.

During those 5 years, he had been in and out of the hospital 5 to 6 times. He had undergone high tech examinations such as angiogram by injecting VINYL via the vein into the heart. He consulted and sought treatment from many good doctors where later he underwent a major surgical operation. Upon his discharge from the hospital, he quit his job to convalesce. However, each time when his illness relapsed, the attack seemed to be even more severe.

Last year, in August, his relatives were in despair and expected he would not live much longer. It so happened at that time that the victim's relative came across antioxidant water processor. His illness responded well and he is now on the road to recovery."

In the United States, cardiovascular diseases account for more than one-half of the approximate 2 million deaths occurring each year.... It is estimated that optimal conditioning of drinking water could reduce this cardiovascular disease mortality rate by as much as 15 percent in the United States

Report of the Safe Drinking Water Committee of the National Academy of Sciences, 1977

Eczema and the Effects of Ionized Water

Prof. Tamura Tatsuji, Keifuku Rehabilitation Center

"Eczema is used to describe several varieties of skin conditions, which have a number of common features. The exact cause or causes of eczema are not fully understood. In many cases, eczema can be attributed by external irritants.

Let me introduce a patient who recovered from skin disease after consuming the antioxidant water. This patient suffered 10 years of eczema and could not be cured effectively even under specialist treatment. This patient, who is 70 years of age, is the president of a vehicle spare parts company. After the war, his lower limbs suffered acute eczema, which later became chronic. He was repeatedly treated in a specialist skin hospital.

The left limb responded well to treatment, but not so on the right limb. He suffered severe itchiness, which, when scratched led to bleeding. During the last 10 years, he was seen and treated by many doctors. When I first examined him, his lower limb around the joints was covered with vesicles. Weeping occurred owing to serum exuding from the vesicles.

I advised him to try consuming antioxidant water. He bought a unit and consumed the antioxidant water religiously and used the acidic water to bathe the affected areas. After 2 weeks of treatment the vesicles dried up. The eczema was completely cleared without any relapse after 1½ month."

Diabetes and the Effects of Ionized Water

Prof. Kuwata Keijiroo, Doctor of Medicine

"When I was serving in the Fire Insurance Association, I used to examine many diabetic patients. Besides treating them with drugs, I provided them with antioxidant water. After drinking antioxidant water for one month, 15 diabetic patients were selected and sent to Tokyo University for further test and observations.

Initially, the more serious patients were a bit apprehensive about the treatment. When the antioxidant water was consumed for some time, the sugar in the blood and urine ranged from a ratio of 300 mg/l to 2 mg / dc. There was a time where the patient had undergone 5 to 6 blood tests a day and detected to be within normal range. Results also showed that even 1 ½ hour after meals, the blood sugar and urine ratio was 100 mg/dc: 0 mg/dc . The sugar in the urine has completely disappeared."

NOTE: More Americans than ever before are suffering from diabetes, with the number of new cases averaging almost 800,000 each year. The disease has steadily increased in the United States since 1980, and in 1998, 16 million Americans were diagnosed with diabetes (10.3 million diagnosed; 5.4 million undiagnosed). Diabetes is the seventh

leading cause of death in the United States , and more than 193,000 died from the disease and its related complication in 1996.

The greatest increase - 76 percent - occurred in people age 30 to 30.

From: U. S. Department of Health and Human Services, October 13, 2000 Fact Sheet.

Digestive Problems and Ionized Water

Prof. Kogure Keizou, Kogure Clinic of Juntendo Hospital

The stomach is readily upset both by diseases affecting the stomach and by other general illnesses. In addition, any nervous tension or anxiety frequently causes gastric upset, vague symptoms when This information is under some strain.

The important role of antioxidant water in our stomach is to neutralize the secretion and strengthen its functions.

Usually, after consuming the antioxidant water for 1 to 3 minutes, the gastric juice increase to 1½ times. For those suffering from hypochlorhydria or achlorhydria (low in gastric juice) the presence of antioxidant water will stimulate the stomach cells to secrete more gastric juice. This in turn enhances digestion and absorption of minerals.

However, on the other hand, those with hyperchlorhydria (high in gastric juice), the antioxidant water neutralizes the excessive gastric juice. Hence, it does not create any adverse reaction.

According to the medical lecturer from Maeba University , the pH of the gastric secretion will still remain normal when antioxidant water is consumed. This proves that the ability of the antioxidant water is able to neutralize as well as to stimulate the secretion."

Reduced Water for the Prevention of Disease

Dr.Sanetaka Shirahata

Graduate school of Genetic Resources Technology , Kyushu University ,

6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581, Japan.

It has long been established that reactive oxygen species (ROS) cause many types of damage to biomolecules and cellular structures that, in turn result in the development of a variety of pathologic states such as diabetes, cancer and aging. Reduced water is defined as anti-oxidative water produced by reduction of water. Electrolyzed reduced water (ERW) has been demonstrated to be hydrogen-rich water and can scavenge ROS in vitro (Shirahata et al., 1997).

The reduction of proton in water to active hydrogen (atomic hydrogen, hydrogen radical) that can scavenge ROS is very easily caused by a weak current, compared to oxidation of hydroxyl ion to oxygen molecule. Activation of water by magnetic field, collision, minerals etc. will also produce reduced water containing active hydrogen and/or hydrogen molecule. Several natural waters such as Hita Tenryosui water drawn from deep underground in Hita city in Japan, Nordenau water in Germany and Tlacote water in Mexico are known to alleviate various diseases.

We have developed a sensitive method by which we can detect active hydrogen existing in reduced water, and have demonstrated that not only ERW but also natural reduced waters described above contain active hydrogen and scavenge ROS in cultured cells. ROS is known to cause reduction of glucose uptake by inhibiting the insulin-signaling pathway in cultured cells. Reduced water scavenged intracellular ROS and stimulated glucose uptake in the presence or absence of insulin in both rat L6 skeletal muscle cells and mouse 3T3/L1 adipocytes. This insulin-like activity of reduced water was inhibited by wortmannin that is specific inhibitor of PI-3 kinase, a key molecule in insulin signaling pathways. Reduced water protected insulin-responsive cells from sugar toxicity and improved the damaged sugar tolerance of type 2 diabetes model mice, suggesting that reduced water may improve insulin-independent diabetes mellitus. Cancer cells are generally exposed to high oxidative stress. Reduced water cause impaired tumor phenotypes of human cancer cells, such as reduced growth rate, morphological changes, reduced colony formation ability in soft agar, passage number-dependent telomere shortening, reduced binding abilities of telomere binding proteins and suppressed metastasis. Reduced water suppressed the growth of cancer cells transplanted into mice, demonstrating their anti-cancer effects in vivo. Reduced water will be applicable to not only medicine but also food industries, agriculture, and manufacturing industries.

Shirahata, S. et al .: Electrolyzed reduced water scavenges active oxygen species and protects DNA from oxidative damage. *Biochem. Biophys. Res. Commun.*, 234, 269174, 1997.

Clinical evaluation of alkaline *Ionized Water* for abdominal complaints: Placebo controlled double blind tests by Hirokazu Tashiro, Tetsuji Hokudo, Hiromi Ono, Yoshihide Fujiyama, Tadao Baba (National Ohkura Hospital, Dept. of Gastroenterology; Institute of Clinical Research, Shiga University of Medical Science, Second Dept. of Internal Medicine)

Effect of alkaline *Ionized Water* on abdominal complaints was evaluated by placebo controlled double blind tests. Overall scores of improvement using alkaline *Ionized Water* marked higher than those of placebo controlled group, and its effect proved to be significantly higher especially in slight symptoms of chronic diarrhea and abdominal complaints in cases of general malaise. Alkaline *Ionized Water* group did not get interrupted in the course of the test, nor did it show serious side effects nor abnormal test data. It was confirmed that alkaline *Ionized Water* is safer and more effective than placebos.

Summary

Effect of alkaline *Ionized Water* on abdominal complaints was clinically examined by double blind tests using clean water as placebo. Overall improvement rate was higher for alkaline *Ionized Water* group than placebo group and the former proved to be significantly more effective than the other especially in cases of slight symptoms. Examining improvement rate for each case of chronic diarrhea, constipation and abdominal complaints, alkaline *Ionized Water* group turned out to be more effective than placebo group for chronic diarrhea, and abdominal complaints. The test was stopped in one case of chronic diarrhea, among placebo group due to exacerbation, whereas alkaline *Ionized Water* group did not stop testing without serious side effects or abnormal test data in all cases. It was confirmed that alkaline *Ionized Water* is more effective than clean water against chronic diarrhea, abdominal complaints and overall improvement rate (relief of abdominal complaints) and safer than clean water.

Introduction

Since the approval of alkaline *Ionized Water* electrolyzers by Pharmaceutical Affairs Law in 1966 for its antacid effect and efficacy against gastrointestinal disorders including hyperchylia, indigestion, abnormal gastrointestinal fermentation and chronic diarrhea, they have been extensively used among patients. However, medical and scientific evaluation of their validity is not established. In our study, we examined clinical effect of alkaline *Ionized Water* on gastrointestinal disorders across many symptoms in various facilities. Particularly, we studied safety and usefulness of alkaline *Ionized Water* by double-blind tests using clean water as a control group.

Test subjects and methods

163 patients (34 men, 129 women, age 21 to 72, average 38.6 years old) of indigestion, abnormal gastrointestinal fermentation (with abnormal gas emission and rugitus) and abdominal complaints caused by irregular dejection (chronic diarrhea, or constipation) were tested as subjects with good informed consent. Placebo controlled double blind tests were conducted using alkaline *Ionized Water* and clean water at multiple facilities. An alkaline *Ionized Water* electrolyzer sold commercially was installed with a pump driven calcium dispenser in each of the subject homes. Tested alkaline *Ionized Water* had pH at 9.5 and calcium concentration at 30ppm. Each subject in placebo group used a water purifier that has the same appearance as the electrolyzer and produces clean water.

The tested equipment was randomly assigned by a controller who scaled off the key code which was stored safely until the tests were completed and the seal was opened again.

Water samples were given to each patient in the amount of 200ml in the morning with the total of 500ml or more per day for a month. Before and after the tests, blood, urine and stool were tested and a log was kept on the subjective symptoms, bowel movements and accessory symptoms. After the tests, the results were analyzed based on the log and the test data.

Test Results

1. Symptom. Among 163 tested subjects, alkaline *Ionized Water* group included 84 and placebo group 79. Background factors such as gender, age and basal disorders did not contribute to significant difference in the results.
2. Overall improvement rate. As to overall improvement rate of abdominal complaints, alkaline *Ionized Water* group had 2 cases of outstanding improvement (2.5%), 26 cases of fair improvement (32.1%), 36 cases of slight improvement (44.4%), 13 cases of no change (16%) and 4 cases of exacerbation (4.9%), whereas placebo group exhibited 4 (5.2%), 19 (24.7%), 27 (35.1%), 25 (32.5%) and 2 cases (2.6%) for the same category. Comparison between alkaline *Ionized Water* and placebo groups did not reveal any significant difference at the level of 5% significance according to the Wilcoxon test, although alkaline *Ionized Water* group turned out to be significantly more effective than placebo group at the level of p value of 0.22.
3. Improvement rate by basal symptom. Basal symptoms were divided into chronic diarrhea, constipation and abdominal complaints (dyspepsia) and overall improvement rate was evaluated for each of them to study effect of alkaline *Ionized Water*. In case of chronic diarrhea, alkaline *Ionized Water* group resulted in 94.1% of effective cases and 5.9% of non effective cases. Placebo group came up with 64.7% effective and 35.3% non effective. These results indicate alkaline *Ionized Water* group proved to be significantly more effective than placebo group. In case of slighter chronic diarrhea, comparison between groups revealed that alkaline *Ionized Water* group is significantly more effective than placebo group (p=0.015). In case of constipation, alkaline *Ionized Water* group consisted of 80.5% of effective

and 19.5% of non effective cases, whereas placebo group resulted in 73.3% effective and 26.3 non effective. As to abdominal complaints (dyspepsia), alkaline *Ionized Water* group had 85.7% of effective and 14.3% non effective cases while placebo group showed 47.1% and 62.9% respectively. Alkaline *Ionized Water* group proved to be significantly more effective than placebo group ($p=0.025$).

Conclusion

As a result of double blind clinical tests of alkaline *Ionized Water* and clean water, alkaline *Ionized Water* was proved to be more effective than clean water against chronic diarrhea, abdominal complaints (dyspepsia) and overall improvement rate (relief from abdominal complaints). Also, safety of alkaline *Ionized Water* was confirmed which clinically verifies its usefulness.

Treatment of Escherichia coli (O157:H7) inoculated alfalfa seeds and sprouts with electrolyzed oxidizing water

International Journal Food Microbiology 2003 Sep 15;86(3):231-7. Department of Agricultural and Biological Engineering, Pennsylvania State University, University Park, PA 16802, USA.

Electrolyzed oxidizing water is a relatively new concept that has been utilized in agriculture, livestock management, medical sterilization, and food sanitation. Electrolyzed oxidizing (EO) water generated by passing sodium chloride solution through an EO water generator was used to treat alfalfa seeds and sprouts inoculated with a five-strain cocktail of nalidixic acid resistant *Escherichia coli* O157:H7. EO water had a pH of 2.6, an oxidation-reduction potential of 1150 mV and about 50 ppm free chlorine. The percentage reduction in bacterial load was determined for reaction times of 2, 4, 8, 16, 32, and 64 min. Mechanical agitation was done while treating the seeds at different time intervals to increase the effectiveness of the treatment. Since *E. coli* O157:H7 was released due to soaking during treatment, the initial counts on seeds and sprouts were determined by soaking the contaminated seeds/sprouts in 0.1% peptone water for a period equivalent to treatment time. The samples were then pummeled in 0.1% peptone water and spread plated on tryptic soy agar with 5 microg/ml of nalidixic acid (TSAN). Results showed that there were reductions between 38.2% and 97.1% (0.22-1.56 log(10) CFU/g) in the bacterial load of treated seeds.

The reductions for sprouts were between 91.1% and 99.8% (1.05-2.72 log(10) CFU/g). An increase in treatment time increased the percentage reduction of *E. coli* O157:H7. However, germination of the treated seeds reduced from 92% to 49% as amperage to make EO water and soaking time increased. EO water did not cause any visible damage to the sprouts.

Ionized Acid Water promotes substantially healthier plant growth.

A clinical study of liver abscesses at the Critical Care and Emergency Center of Wate Medical University]

Fujino Y, Inoue Y, Onodera M, Yaegashi Y, Sato N, Endo S, Omori H, Suzuki K. [Article in Japanese] Department of Critical Care Medicine, Wate Medical University.

We studied 13 emergency cases of liver abscess. Five cases of septic shock or clouding of consciousness were identified on admission. Six patients had diabetes mellitus. Twelve patients met the diagnostic criteria for systemic inflammatory response syndrome, and nine met the criteria for disseminated intravascular coagulation. Plasma endotoxin levels improved rapidly after drainage. Causative organisms were isolated in all patients, and the most common organism was *Klebsiella pneumoniae* (seven cases). Percutaneous transhepatic abscess drainage (PTAD) was performed not only in single cases but also in multiple cases with main huge abscesses. Surgical treatment was performed in the following three cases: a ruptured abscess, an ineffective PTAD, and a case of peritonitis after PTAD. Irrigation of abscesses with strong acidic electrolyzed water revealed a significant decrease in treatment duration. In the majority of our cases, severe conditions were identified on admission. Strong acidic electrolyzed water was useful for management of PTAD.

Evaluation of hypochlorous acid washes in the treatment of chronic venous leg ulcers. *J.B. Selkon, MBChB, FRCPath, DCP, Honorary Consultant Microbiologist, Oxford University, Department of Clinical Medicine; G.W. Cherry, DPhil (Oxon), Clinical Faculty Member, Oxford University Medical School; Chairman, Oxford International Wound Healing Foundation; J.M. Wilson, RN, Clinical Research Nurse; M.A. Hughes, PhD, Clinical Scientist, Oxford International Wound Healing Foundation; all at Dermatology Department, Churchill Hospital, Oxford, UK. Email: jselkon@onetel.com* VOL 15, NO 1, Page 33, January 2006

Hypochlorous acid is a highly microbial agent active against bacteria, viruses and fungi. This study aimed to determine if it has a role as an additional treatment for chronic venous leg ulcers that have not healed with standard treatment.

The patients acted as their own controls, in that only those who failed to achieve a 44% reduction in wound size with compression therapy after three weeks received the new treatment. Of the 30 patients recruited into the study, 20 were given the hypochlorous acid washes. In these patients, nine ulcers healed and five reduced in size by 60% in the follow-up period. All patients' ulcers became pain-free.

Enhancing the bactericidal effect of electrolyzed water on *Listeria monocytogenes* biofilms formed on stainless steel. Ayebah B, Hung YC, Frank JF. Department of Food Science and Technology, University of Georgia, 1109 Experiment Street, Griffin, Georgia 30223, USA.

Biofilms are potential sources of contamination to food in processing plants, because they frequently survive sanitizer treatments during cleaning. The objective of this research was to investigate the combined use of alkaline and acidic electrolyzed (EO) water in the inactivation of *Listeria monocytogenes* biofilms on stainless steel surfaces. Biofilms were grown on rectangular stainless steel (type 304, no. 4 finish) coupons (2 by 5 cm) in a 1:10 dilution of tryptic soy broth that contained a five-strain mixture of *L. monocytogenes* for 48 h at 25 degrees C. The coupons with biofilms were then treated with acidic EO water or alkaline EO water or with alkaline EO water followed by acidic EO water produced at 14 and 20 A for 30, 60, and 120 s. Alkaline EO water alone did not produce significant reductions in *L. monocytogenes* biofilms when compared with the control. Treatment with acidic EO water only for 30 to 120 s, on the other hand, reduced the viable bacterial populations in the biofilms by 4.3 to 5.2 log CFU per coupon, whereas the combined treatment of alkaline EO water followed by acidic EO water produced an additional 0.3- to 1.2-log CFU per coupon reduction. The population of *L. monocytogenes* reduced by treatments with acidic EO water increased significantly with increasing time of exposure. However, no significant differences occurred between treatments with EO water produced at 14 and 20 A. Results suggest that alkaline and acidic EO water can be used together to achieve a better inactivation of biofilms than when applied individually.

Selective stimulation of the growth of anaerobic microflora in the human intestinal tract by electrolyzed reducing water. Vorobjeva NV. Department of Physiology of Microorganisms, Biology Faculty, Lomonosov Moscow State University, 119992 Moscow, Russia. nvvorobjeva@mail.ru

96-99% of the "friendly" or residential microflora of intestinal tract of humans consists of strict anaerobes and only 1-4% of aerobes. Many diseases of the intestine are due to a disturbance in the balance of the microorganisms inhabiting the gut. The treatment of such diseases involves the restoration of the quantity and/or balance of residential microflora in the intestinal tract. It is known that aerobes and anaerobes grow at different oxidation-reduction potentials (ORP). The former require positive E(h) values up to +400 mV. Anaerobes do not grow unless the E(h) value is negative between -300 and -400 mV. In this work, it is suggested that prerequisite for the recovery and maintenance of obligatory anaerobic microflora in the intestinal tract is a negative ORP value of the intestinal milieu. Electrolyzed reducing water with E(h) values between 0 and -300 mV produced in electrolysis devices possesses this property. Drinking such water favors the growth of residential microflora in the gut. A sufficient array of data confirms this idea. However, most researchers explain the mechanism of its action by an antioxidant properties destined to detox the oxidants in the gut and other host tissues. Evidence is presented in favor of the hypothesis that the primary target for electrolyzed reducing water is the residential microflora in the gut.

Efficacy of electrolyzed water in inactivating *Salmonella enteritidis* and *Listeria monocytogenes* on shell eggs. Journal of Food Protection 2005 May;68(5):986-90. Park CM, Hung YC, Lin CS, Brackett RE. Department of Food Science and Technology, University of Georgia, Griffin, Georgia 30223-1797, USA.

The efficacy of acidic electrolyzed (EO) water produced at three levels of total available chlorine (16, 41, and 77 mg/liter) and chlorinated water with 45 and 200 mg/liter of residual chlorine was investigated for inactivating *Salmonella Enteritidis* and *Listeria monocytogenes* on shell eggs. An increasing reduction in *Listeria* population was observed with increasing chlorine concentration from 16 to 77 mg/liter and treatment time from 1 to 5 min, resulting in a maximal reduction of 3.70 log CFU per shell egg compared with a de-ionized water wash for 5 min. There was no significant difference in antibacterial activities against *Salmonella* and *Listeria* at the same treatment time between 45 mg/liter of chlorinated water and 14-A acidic EO water treatment ($P > \text{or} = 0.05$). Chlorinated water (200 mg/liter) wash for 3 and 5 min was the most effective treatment; it reduced mean populations of *Listeria* and *Salmonella* on

inoculated eggs by 4.89 and 3.83 log CFU/shell egg, respectively. However, reductions (log CFU/shell egg) of *Listeria* (4.39) and *Salmonella* (3.66) by 1-min alkaline EO water treatment followed by another 1 min of 14-A acidic EO water (41 mg/liter chlorine) treatment had a similar reduction to the 1-min 200 mg/liter chlorinated water treatment for *Listeria* (4.01) and *Salmonella* (3.81). This study demonstrated that a combination of alkaline and acidic EO water wash is equivalent to 200 mg/liter of chlorinated water wash for reducing populations of *Salmonella* Enteritidis and *L. monocytogenes* on shell eggs.

Science Direct-Application of electrolyzed oxidizing water to reduce *Listeria monocytogenes* on ready-to-eat meats. Friday, April 1st, 2005 K.A. Fabrizio, and C.N. Cutter, Department of Food Science, 111 Borland Laboratory, The Pennsylvania State University, University Park, PA 16802, USA

Experiments were conducted to determine the effectiveness of acidic (EOA) or basic electrolyzed oxidizing (EOB) water, alone or in combination, on ready-to-eat (RTE) meats to reduce *Listeria monocytogenes* (LM). Frankfurters or ham surfaces were experimentally inoculated with LM and subjected to dipping or spraying treatments (25 or 4 °C for up to 30 min) with EOA, EOB, and other food grade compounds. LM was reduced the greatest when frankfurters were treated with EOA and dipped at 25 °C for 15 min. A combination spray application of EOB/EOA also resulted in a slight reduction of LM on frankfurters and ham. However, reductions greater than 1 log CFU/g were not observed for the duration of the study. Even with a prolonged contact time, treatments with EOA or EOB were not enough to meet regulatory requirements for control of LM on RTE meats. As such, additional studies to identify food grade antimicrobials to control the pathogen on RTE meats are warranted

CONTROL OF POWDERY MILDEW BY SPRAYING THE ELECTROLYZED WATER IN HYDROPONICALLY GROWN STRAWBERRY ISHS Acta Horticulturae 559: V International Symposium on Protected Cultivation in Mild Winter Climates: Current Trends for Sustainable Technologies

Abstract:

Effect of spraying electrolyzed acid water and alkaline water obtained by electrolysis of KCl solution on the incidence of powdery mildew (*Sphaerotheca macularis*) is investigated in strawberry (*Fragaria x ananassa* Duch. cv. Nyoho) grown in hydroponics. Free effective chlorine concentration of the acid water at 0.5m from spraying nozzle ranged between 5.7 and 15.6 mg/l. pH and oxidation-reduction potential of the acid water ranged from 2.2 to 2.3 and from 1100 to 1140 mV, and those of the alkaline water ranged from 12.3 to 12.6 and from 30 to 280 mV, respectively. The treatment plots were 1. No spraying (NSp), 2. Acid water spraying (Ac), 3. Alkaline water spraying 30 minutes after acid water spraying (Ac+Al), and 4. Agricultural chemicals spraying (Chem). Those electrolyzed waters were sprayed onto the leaves and petioles once every week, and chemicals were sprayed once every two or three weeks. The results indicate that the acid water spraying and acid + alkaline water spraying can control the powdery mildew in strawberry and can reduce the use of chemical fungicide in protected cultivation.

Science Direct-Toxicity of mixed-oxidant electrolyzed oxidizing water to in vitro and leaf surface populations of vegetable bacterial pathogens and control of bacterial diseases in the greenhouse. Tuesday, December 28th, 2004. Ken Pernezny, , Richard N. Raid, Nikol Havranek and Jairo Sanchez. IFAS, Everglades Research and Education Center, University of Florida, 3200 E. Palm Beach Road, Belle Glade, FL 33430, USA

Efficacy of electrolyzed oxidizing water for the microbial safety and quality of eggs. Poultry Science 2004 Dec;83(12):2071-8. Bialka KL, Demirci A, Knabel SJ, Patterson PH, Puri VM. Department of Agricultural & Biological Engineering, The Pennsylvania State University, University Park, Pennsylvania 16802, USA. During commercial processing, eggs are washed in an alkaline detergent and then rinsed with chlorine to reduce dirt, debris, and microorganism levels. The alkaline and acidic fractions of electrolyzed oxidizing (EO) water have the ability to fit into the 2-step commercial egg washing process easily if proven to be effective. Therefore, the efficacy of EO water to decontaminate *Salmonella* Enteritidis and *Escherichia coli* K12 on artificially inoculated shell eggs was investigated. For the in vitro study, eggs were soaked in alkaline EO water followed by soaking in acidic EO water at various temperatures and times. Treated eggs showed a reduction in population between $> \text{or} = 0.6$ to $> \text{or} = 2.6 \log_{10}$ cfu/g of shell for *S. Enteritidis* and $> \text{or} = 0.9$ and $> \text{or} = 2.6 \log_{10}$ for *E. coli* K12. Log₁₀ reductions of 1.7 and 2.0 for *S. Enteritidis* and *E. coli* K12, respectively, were observed for typical commercial detergent-sanitizer treatments, whereas log₁₀ reductions of $> \text{or} = 2.1$ and $> \text{or} = 2.3$ for *S. Enteritidis* and *E. coli* K12, respectively, were achieved using the EO

water treatment. For the pilot-scale study, both fractions of EO water were compared with the detergent-sanitizer treatment using E. coli K12. Log10 reductions of $> \text{ or } = 2.98$ and $> \text{ or } = 2.91$ were found using the EO water treatment and the detergent-sanitizer treatment, respectively. The effects of 2 treatments on egg quality were investigated. EO water and the detergent-sanitizer treatments did not significantly affect albumen height or eggshell strength; however, there were significant affects on cuticle presence. These results indicate that EO water has the potential to be used as a sanitizing agent for the egg washing process.

Susceptibility of *Penicillium expansum* spores to sodium hypochlorite, electrolyzed oxidizing water, and chlorine dioxide solutions modified with nonionic surfactants. *Journal of Food Protection*. 2006 Aug;69(8):1944-8. Okull DO, Demirci A, Rosenberger D, LaBorde LF.

The disinfection of the surgery department of a garrison hospital using a neutral anolyte. *Voen Med Zh*. 1999 Sep;320(9):56-8. Chueva IM, Mistriukov VV, Mikhailov SI. [Article in Russian]

Methods and resources of sterilization and disinfection in medicine. Abstracts (28 September - 2 October 1992). Russian Conference.

Electrochemistry. *Electrochemistry and Medicine*. // *Itogi nauki i tekhniki*. VINITI. Ed. Polukarov YM, vol. 31, 1990. Article in Russian.

Other Books that Contain References about Ionized Water:

Handbook of Drinking Water Quality, by John De Zuane. Publisher: John Wiley and Sons. 1997. Pages: 592

Pain Free With Far Infrared Mineral Therapy: The Miracle Lamp, by Kara Lee Schoonover. Publisher: iUniverse ISBN: 0595272630. 2003.

Sensual for Life: The Natural Way to Maintain Sexual Vitality, by George L. Redmon. Publisher: Kensington Books ISBN: 0758201389. 224 pgs. 2003.

Natural Choices for Fibromyalgia: Discover Your Personal Method for Pain Relief, by N. D. Oelke, Jane Oelke. Publisher: Natural Choices, Inc. ISBN: 0971551200. 2001. 146 pgs.

Light Through Food: A Reference Guide to Thriving in the Energy of the Fourth Dimension, by Donna Boynton (Channeler), Donna Boynton, Angela McLean. Publisher: Trafford Publishing ISBN: 1412068495. 2005. 100 pgs.

Qi Energy for Health and Healing: A Comprehensive Guide to Accessing Your Healing Energy, by Mallory Fromm. Publisher: Avery. ISBN: 1583331573. 2003. 219 pgs.

"Ionized Water is also very helpful for alleviating the side effects of anesthesia."

Romo: My Life on the Edge, by Bill Romanowski, Published by William Morrow. ISBN: 006075866X. Publisher: HarperTorch. 2005.

A Complete Guide To Chi-Gung, By Daniel Reid. ISBN: 1570625433 Publisher: Shambhala. 2000. 326 pgs.

Secrets of an Alkaline Body. The New Science of Colloidal Biology, by Annie Padden Jubb and David Jubb. Publisher: North Atlantic Books. ISBN: 1556434812. 275 pgs. 2004.

Timeless Secrets of Health and Rejuvenation, By Andreas Moritz. ISBN: 097657151X; pg. 496. 2005. Publisher: Ener-Chi Wellness Center, LLC

Reverse Aging, by Sang Whang. ISBN: 0966236327, Published by JSP Publishing. 2001.

Fantastic Voyage: Live Long Enough to Live, by Ray Kurzweil, Terry Grossman **ISBN:** 0452286670. 464 pages.

ⁱ The cavity in which the large intestine begins and into which the ileum opens. The large blind pouch forming the beginning of the large intestine. Also called *blind gut*.